arXiv:math/0602404v2 [math.DG] 22 Feb 2006

Conformal β- change in Finsler spaces

S. H. Abed [*]

Abstract. We investigate what we call a conformal β - change in Finsler spaces, namely

$$L(x,y) \to {}^*L(x,y) = e^{\sigma(x)}L(x,y) + \beta(x,y)$$

where σ is a function of x *only and* $\beta(x,y)$ is a given 1- form.

This change generalizes various types of changes: conformal changes, Randers changes and β - changes.

Under this change, we obtain the relationships between some tensors associated with (M, L) and the corresponding tensors associated with $(M, {}^*L)$. We investigate some σ-invariant tensors . This investigation allows us to give an answer to the question: Are the properties of C-reducibility, S_3-likeness and S_4-likeness invariant under a conformal β - change?

1. Introduction and Notations

Let (M, L) be a Finsler space, where M is an n-dimensional differentiable manifold equipped with a fundamental function L. Given a function σ, the change

$$L(x,y) \longrightarrow e^{\sigma(x)}L(x,y)$$

is called a conformal change. The conformal theory of Finsler spaces has been initiated by M.S. Kneblman [5] in 1929 and has been deeply investigated by many authors: [1], [3], [4],... etc

In 1941, Randers [9] has introduced the Finsler change

$$^{r}L(x,y) \longrightarrow {}^{r}L(x,y) + \beta(x,y)$$

where ^{r}L is a Riemanian structur and β is a 1-form on M. The resulting space is a Finsler space. This change has been studied by several authors: [7],[12],...etc.

The Randers change has been generalized by Shibata [10] to what is called a β - change

$$L(x,y) \longrightarrow L(x,y) + \beta(x,y)$$

---

[*] Department of Mathematics, Faculty of Science, Cairo University, Cairo, Egypt.

where L a fundamental Finslerian function. The resulting space known as a generalized Randers space was studied in [13], [4], [7], [11] and [8],... etc.

In this paper, we construct a theory which generalizes all the above mentioned changes. In fact, we consider a change of the form

$$L(x,y) \longrightarrow {}^*L(x,y) = e^{\sigma(x)}L(x,y) + \beta(x,y),$$

where σ is a function of x and $\beta(x,y) = b_i(x)y^i$ is a 1- form on M, which we call a conformal β - change. This change generalizes various type of changes. When $\beta = 0$, it reduces to a conformal change. When $\sigma = 0$, it reduces to a β- change and consequently to a Randers change.

We obtain the relationships between some tensors associated with (M, L) (the fundamental tensor, the $h(hv)$ - torsion and the third curvature tensor) and the corresponding tensors associated with $(M, {}^* L)$.

Under the conformal β - change, we investigate some σ- invariant tensors (a tensor K is σ- invariant if ${}^*K(x,y) = e^{\sigma}K(x,y)$).

This investigation leads us to find out necessary and /or sufficient conditions for the properties of C-reducibility, S_3-likeness and S_4-likeness to be invariant under a conformal β - change (cf. theorems A, B, and C).

More investigation and development of this theory will be the object of forthcoming papers.

Throughout the present paper, (x^i) denotes the coordinates of a point of the base manifold M and (y^i) the supporting element $(\dot{x}^i)$.

We use the following notations:

$l_i := \dot{\partial}_i L \ = \frac{\partial L}{\partial y^i}$: the normalized supporting element,
$h_{ij} := L\dot{\partial}_i l_i = Ll_{ij}$: the angular metric tensor,
$g_{ij} := \frac{1}{2}\dot{\partial}_i \ \dot{\partial}_j L^2$: the fundamental tensor,
$c_{ijk} := \ \dot{\partial}_k\,(g_{ij}/2)$: the(h) hv -torsion tensor ,
$c_i := g^{jk}\,c_{ijk}$: the torsion vector,
$c^k := g^{jk}\,c_i$: $\ , c^2 = c_i c^i$,

$S_{hijk} := c_{ijr}c_{h\ k}^{\ r} - c_{ikr}c_{h\ j}^{\ r}$: the components of the third curvature tensor.

2. Conformal β-change

We firstly introduce the following definition

Definition 1. *A change of Finsler metric defined by*

$$L(x,y) \longrightarrow {}^*L(x,y) = e^{\sigma(x)}L(x,y) + \beta(x,y) \tag{1}$$

where $\sigma = \sigma(x)$ is a function of x and $\beta(x,y) = b_i(x)y^i$ is a 1- form, will be called a conformal β-change.

This change generalizes various changes studied by Randers [9], Matsumuto [7], Shibata.[10]...etc.

We assume that ${}^*L(x,y)$ enjoys the same properties possessed by $L(x,y)$.

As the Finsler space associated to L is denoted by (M,L), we denote the Finsler space associated to the conformal $\beta-$change by $(M,{}^*L)$.

Throughout the whole paper, the geometric objects associated with ${}^*L(x,y)$ will be asterisked.

Definition 2. *A geometric object K is said to be $\sigma-$invariant if it is invariant, up to a factor $e^{\sigma(x)}$, under a conformal $\beta-$change: ${}^*K = e^{\sigma(x)}K$.*

It follows from (1) that

$$ {}^*l_i(x,y) = e^{\sigma(x)}\, l_i(x,y) + b_i(x), \;\; {}^*l_{ij}(x,y) = e^{\sigma(x)}\, l_{ij}(x,y). \tag{2}$$

The angular metric tensor h_{ij} is given in terms of h_{ij} by

$$ {}^*h_{ij} \;=\; {}^*L\,{}^*l_{ij} \;=\; {}^*Le^{\sigma}l_{ij} \;= \tau h_{ij}, \quad \tau = e^{\sigma(x)}\frac{{}^*L}{L}. \tag{3}$$

Then we have the following

Lemma 1. *$\frac{h_{ij}}{L}$ is $\sigma-$ invariant under a conformal $\beta-$change :*

$$ \frac{{}^*h_{ij}}{{}^*L} = e^{\sigma}\,\frac{h_{ij}}{L}. \tag{4}$$

As $h_{ij} = g_{ij} - l_i l_j$, equations (3) give us a relation between the fundamental tensors g_{ij} and ${}^*g_{ij}$:

$$ {}^*g_{ij} = \;\tau(g_{ij} - l_i l_j) + {}^*l_i\,{}^*l_j. \tag{5}$$

The relation between the corresponding covariant components is obtained in the form

$$ {}^*g^{ij} = \;\tau^{-1}g^{ij} + \mu\, l^i\, l^j - \tau^{-2}(\, l^i b^j + l^j b^i), \tag{6}$$

where $\mu = (e^{\sigma}Lb^2 + \beta\,)/\,{}^*L\tau^2$, $\quad b^2 = b_i b^i$, $b^i = g^{ij}b_j$.

Let us introduce the $\pi-$vector field

$$ \overline{m} = \overline{B} - \frac{\beta}{L^2}\overline{\eta}\,, \;\; m^i = b^i - \frac{\beta}{L}l^i, \;\; m^2 = m_i\, m^i.$$

Lemma 2. *For a conformal $\beta-$change which is not conformal (i,e $\beta \neq 0$), $\overline{m} \neq 0$.*

In fact , if $\overline{m} = 0$, then $m_i = 0$ for all i, and consequently $\mathrm{b}_i = \frac{\beta}{L} l_i$ which implies $\beta = e^{\psi(x)} L$, for some function $\psi(x)$.

Lemma 3. *The(h) hv- torsion tensor ${}^*c_{ijk}$ associated to ${}^*F$ is given by*

$$ {}^*c_{ijk} = \tau[c_{ijk} + \frac{1}{2{}^*L} h_{ijk}], \tag{7} $$

where

$$ h_{ijk} = h_{ij} m_k + h_{jk} m_i + h_{ki} m_j. \tag{8} $$

From the tensor ${}^*c_{ijk}$, we obtain the following important tensors:

$$ {}^*c_i{}^r{}_j = c_i{}^r{}_j + \frac{1}{2{}^*L}(h_{ij} m^r + h_j{}^r m_i + h^r{}_i m_j) - \tau^{-1}\, c_{ijs} l^r b^s - \frac{1}{2{}^*L\tau}(2 m_i m_j + m^2 h_{ij}) l^r, \tag{9} $$

$$ {}^*c_i = c_i + \frac{n+1}{2{}^*L} m_i, \tag{10} $$

$$ {}^*c^k = \tau^{-2}[\tau c^k - c_\beta l^k + \frac{n+1}{2{}^*L}(\tau m^k - m^2 l^k)], \tag{11} $$

$$ {}^*c^2 = \tau^{-1}[c^2 + \frac{n+1}{{}^*L}\, A_\beta], \tag{12} $$

$$ {}^*c_\beta = c_\beta + \frac{n+1}{2{}^*L} m^2 \tag{13} $$

where $A_\beta = c_\beta + \frac{n+1}{4{}^*L} m^2, \quad c_\beta = c_i b^i$.

Proof.

- Equation(7) is deduced from the definition of c_{ijk} together with (5)
- Equation(9) is deduced by rasing the index k in (7), using (6)
- Equation(10) is obtained by contracting the subscript i and the superscript r in (9)
- Equation(11) follows from (10) by rasing its subscript, using (6)
- Equation(12) follows directly from (10) and (11) by contracted multiplication
- Equation(13) is obtained easily from (10) and the definition of c_β by contracted multiplication. □

Lemma 4.

(a) *The relation between* ${}^{*}S_{hijk}$ *and* S_{hijk} *takes the form*

$$ {}^{*}S_{hijk} = \tau S_{hijk} - \frac{\tau}{2\,{}^{*}L}\left[h_{ik}H_{jh} + h_{jh}H_{ik} - h_{hk}H_{ij} - h_{ij}H_{hk}\right], \qquad (14) $$

where

$$ H_{ij} = c_i{}^r{}_j m_r + \frac{1}{2\,{}^{*}L} m_i m_j + \frac{1}{4\,{}^{*}L} h_{ij} m^2. \qquad (15) $$

(b) *The* $v-$ *Ricci tensor* ${}^{*}S_{ik}$ *is written in the form*

$$ {}^{*}S_{ik} = S_{ik} - \frac{1}{2\,{}^{*}L}[A_\beta h_{ik} + (n-3)H_{ik}] \qquad (16) $$

(c) *The* $v-$ *scaler curvature tensor is written in the form*

$$ {}^{*}S = \tau^{-1}[S - \frac{n-2}{{}^{*}L} A_\beta] \qquad (17) $$

Proof.

(a) From equations (7) and (9) we have

$$ \begin{aligned} {}^{*}c_{ijr}\ {}^{*}c_h{}^r{}_k &= \tau[c_{ijr} + \frac{1}{2\,{}^{*}L} h_{ijr}][c_h{}^r{}_k + \frac{1}{2\,{}^{*}L}(h_{hk}m^r + h_h{}^r m_k + h^r{}_h m_k) \\ &- \tau^{-1}\, c_{hks} l^r b^s - \frac{1}{2\,{}^{*}L\,\tau}(2m_h m_k + m^2 h_{hk}) l^r] \\ &= \tau c_{ijr}\ c_h{}^r{}_k \ + \frac{\tau}{2\,{}^{*}L}[(c_i{}^r{}_j\ h_{hk} + c_h{}^r{}_k\ h_{ij})\ m_r \\ &+ (c_{ijk}\ m_h\ +\ c_{jkh}\ m_i +\ c_{khi}\ m_j\ +\ c_{hij}\ m_k)\ ] \\ &+ \frac{\tau}{4\,{}^{*}L^2}[h_{ij}\ h_{hk}\ \ m^2 + 2h_{hk}\ m_i\ m_j + 2h_{ij}\ m_h\ m_k\ + h_{jh}\ m_i\ m_k \\ &+ h_{jk} m_i m_h + h_{ih} m_j m_k + h_{ik}\ m_j\ m_h]. \end{aligned} $$

Similarly, one can obtain ${}^{*}c_{ikr}\ {}^{*}c_h{}^r{}_j$ (by interchange j and k). Hence the result.

(b) follows from (a) by contracted multiplication, using (6).

(c) is obtained from (b), using (6) again, by contracted multiplication. □

Remark . *The tensor* H_{ij} *defined by (15) has the properties:*

1. H_{ij} *is a symmetric tensor :* $H_{ij} = H_{ji}$,
2. H_{ij} *is an indicatory tensor :* $H_{ij}\, y^i = 0 = H_{ij}\, y^j$,
3. $g^{ij} H_{ij} = A_\beta$.

3. Geometrical properties of the conformal β-change

Definition 3. *[6] A Finsler space (M, L) of dimension $n \geq 3$ is called a C-reducible space if the h(hv)-torsion tensor c_{ijk} has the form*

$$c_{ijk} = h_{ij}M_k + h_{kj}M_i + h_{ki}M_j \ , \quad M_i = \frac{c_i}{n+1}. \tag{18}$$

Define the tensor

$$K_{ijk} = [c_{ijk} - (h_{ij}M_k + h_{kj}M_i + h_{ki}M_j)]/L.$$

It is clear that K_{ijk} is a symmetric and indicatory tensor. Moreover K_{ijk} vanishes if and only if the Finsler space is C-reducible.

Proposition 1. *Under a conformal β-change, the tensor K_{ijk} is $\sigma-$invariant:*

$${}^*K_{ijk} = e^{\sigma} K_{ijk}.$$

Proof. Using Equation (7) together with the definition of K_{ijk},we get

$$\begin{aligned}
{}^*K_{ijk} &= [{}^*c_{ijk} - ({}^*h_{ij}\ {}^*M_k +^* h_{ij}\ {}^*M_i +^* h_{ki}\ {}^*M_j)]\,/{}^*L \\
&= \tau[(c_{ijk} + \frac{1}{2{}^*L}h_{ijk}) - ({}^*h_{ij}\ {}^*M_k +^* h_{ij}\ {}^*M_i +^* h_{ki}\ {}^*M_j)]\,/{}^*L \\
&= \tau[c_{ijk} + \frac{1}{2{}^*L}(h_{ij}m_k + h_{jk}m_i + h_{ki}m_j.) - \frac{1}{n\ +1}(h_{ij}\ {}^*c_k + h_{jk}\ {}^*c_i + h_{ki}\ {}^*c_j)]\,/{}^*L \\
&= \tau[c_{ijk} + \frac{1}{n\ +1}(h_{ij}\ c_k + h_{jk}c_i + h_{ki}c_j.)]\ /{}^*L \\
&= e^{\sigma}[\ c_{ijk} - (h_{ij}M_k + h_{kj}M_i + h_{ki}M_j)]\,/L = e^{\sigma}K_{ijk}\ . \ \square
\end{aligned}$$

Now, Proposition 1 yields

Theorem A. *Under a conformal β-change $L \longrightarrow {}^*L$, the space (M, L) is C-reducible if and only if the space $(M,^* L)$ is C-reducible.*

Consequently the C-reducibility property is invariant under this change.

It shout be noticed that Theorem 4-1 and Corollary 4-1 of Shibata [10] result from the above Theorem as a very special case. Some results of Matsumoto [7] are also contained in the above Theorem.

Definition 4. *[2] A Finsler space (M, L) of dimension $n > 4$ is called an S_4-like space if the vertical curvature tensor S_{hijk} has the form*

$$S_{hijk} = h_{jh}M_{ik} + h_{ik}M_{jh} - h_{hk}M_{ij} - h_{ij}M_{hk}, \tag{19}$$

where M_{ij} is the symmetric and indicatory tensor given by $M_{ij} = \frac{1}{n-3}[S_{ij} - \frac{Sh_{ij}}{2(n-2)}]$.

Define the tensor

$$\eta_{hijk} = [S_{hijk} - (h_{jh}M_{ik} + h_{ik}M_{jh} - h_{hk}M_{ij} - h_{ij}M_{hk})]/L,$$

It is clear that η_{hijk} vanishes if and only if the manifold (M, L) is an S_4-like manifold.

It is not difficult to prove the following.

Lemma 5. *The tensor* $^*M_{ij}$ *is given in termes of* M_{ij} *by*

$$^*M_{ij} = M_{ij} \ - \frac{1}{2^*L}H_{ij}.$$

In fact, the result follows from (12) and (16).

Proposition 2. *Under a conformal* β*-change, the tensor* η_{hijk} *is* $\sigma-$*invariant :*

$$^*\eta_{hijk} = e^{\sigma}\eta_{hijk}$$

Proof. Taking Lemma 4a and Lemma 5 into account, we get

$$\begin{aligned}
^*L^*\eta_{hijk} &= {}^*S_{hijk} - ({}^*h_{jh}{}^*M_{ik} +^* h_{ik}{}^* M_{jh} -^* h_{hk}{}^*M_{ij} -^* h_{ij}{}^*M_{hk}) \\
&= \tau S_{hijk} - \frac{\tau}{2^*L}[h_{jk}H_{ih} + h_{ih}H_{jk} - h_{hk}H_{ij} - h_{ij}H_{hk}] \\
&- \tau[h_{jk}(M_{ih} - \frac{1}{2^*L}H_{ih}) + h_{ik}(M_{jh} - \frac{1}{2^*L}H_{jh}) \\
&- h_{hk}(M_{ij} - \frac{1}{2^*L}H_{ij}) - h_{ij}\ (M_{hk} - \frac{1}{2^*L}H_{hk})] \\
&= \tau[S_{hijk} - (h_{jk}M_{ih} + h_{ih}M_{jk} - h_{hk}M_{ij} - h_{ij}M_{hk})] \\
&= \tau\ L\ \eta_{hijk} = \frac{e^{\sigma}\ {}^*L}{L}\ L\ \eta_{hijk} = e^{\sigma}\ {}^*L\ \eta_{hijk}.
\end{aligned}$$

Hence the result. □

Proposition (2) yields

Theorem B. *Under a conformal* β*-change* $L \longrightarrow {}^*L$*, the space* (M, L) *is* S_4-*like if and only if the space* $(M,^* L)$ *is an* S_4-*like.*

Consequently, the S_4-*likeness property is invariant under this change.*

The above result generalizes Theorem 4-5 (and its Corollary) of Shibata [10] .

Definition 5. *[2] A Finsler space* (M, L) *of dimension* $n > 3$ *is called an* S_3-*like space if the vertical curvature tensor* S_{hijk} *has the form*

$$S_{hijk} = \frac{S}{(n-1)\ (n-2)}\,[h_{ik}h_{jh} - h_{ij}h_{hk}] \qquad (20)$$

Define the tensor

$$\zeta_{hijk} \;= [\; S_{hijk} - \frac{S}{(n-1)\;(n-2)}\,(h_{ik}h_{jh} - h_{ij}h_{hk})]\; lL$$

It is clear that ζ_{hijk} vanishes if and only if the manifold (M, L) is an S_3-like manifold

Proposition 3. *Under a conformal β-change, the tensor ζ_{hijk} is $\sigma-$invariant if and only if $H_{ij} = \frac{1}{n-1}A_\beta h_{ij}$.*

Proof. Using Equation (7) together with the definition of K_{ijk}, we get

$$\begin{aligned}
{}^*L{}^*\zeta_{hijk} &= [{}^*S_{hijk} - \frac{{}^*S}{(n-1)\;(n-2)}\,({}^*h_{ik}\;{}^*h_{jh} -{}^*\, h_{ij}\;{}^*h_{hk})] \\
&= [\tau S_{hijk} - \frac{\tau}{2\;{}^*L}(h_{ik}\,H_{jh} + h_{jh}H_{ik} - h_{hk}H_{ij} - h_{ij}H_{hk}) \\
&- \frac{\tau}{(n-1)\;(n-2)}\,(S - \frac{(n-2)}{{}^*L}A_\beta)(h_{ik}\;h_{jh} - h_{ij}\;h_{hk})] \\
&= [\tau(S_{hijk} - \frac{S}{(n-1)\;(n-2)}\,(h_{ik}h_{jh} - h_{ij}h_{hk}))] \\
&- \frac{\tau}{2\;{}^*L}[(h_{ik}\,H_{jh} + h_{jh}H_{ik} - h_{hk}H_{ij} - h_{ij}H_{hk}) + \frac{1}{(n-1)}A_\beta(2h_{ik}\;h_{jh} - 2h_{ij}\;h_{hk})] \\
&= \tau L\;\zeta_{hijk}\; - \frac{\tau}{2\;{}^*L}[\,(h_{ik}\,(H_{jh} - \frac{1}{(n-1)}A_\beta h_{jh})) + (h_{jh}\,(H_{ik} - \frac{1}{(n-1)}A_\beta h_{ik})) + \\
&+ \;(h_{hk}\,(H_{ij} - \frac{1}{(n-1)}A_\beta h_{ij})) + h_{ij}(H_{hk} - \frac{1}{(n-1)}A_\beta h_{hk}))]
\end{aligned}$$

Now the tensor ζ^*_{hijk} is σ-invariant (${}^*\zeta_{hijk} = e^\sigma\zeta_{hijk}$) if and only if all terms of the forms $H_{ij} - \frac{1}{(n-1)}A_\beta h_{ij}$ vanish; that is, if and only if the condition $H_{ij} = \frac{1}{n-1}A_\beta h_{ij}$ holds. □
Consequently we get

Theorem C. *Under a conformal β-change $L \longrightarrow {}^*L$, the following two assertions are equivalent*

1. *The space (M, L) is S_3-like ,*
2. *The space $(M,{}^* L)$ is S_3-like*

if and only if the condition $H_{ij} = \frac{1}{n-1}A_\beta h_{ij}$ holds.

Consequently, the S_3- likeness property is invariant under this change if and only if $H_{ij} = \frac{1}{n-1}A_\beta h_{ij}$

References

[1] M. Hashiguchi, On conformal transformation of Finsler metric, J. Math. Kyoto Univ.. 16(1976) pp. 25-50.

[2] F. Ikedo, On S_3-and S_4-like Finsler spaces with the T- tensor of a special form, Tensor, N.S..35(1981) pp. 345-351.

[3] H. Izumi, Conformal transformations of Finsler spaces I and II, Tensor, N.S.. 31 and 33(1977 and 1980) pp. 33-41 and 337-359.

[4] M. Kitayama, Geometry of transformations of Finsler metrics, Hokkaido University of Education, Kushiro Compus.. Japan(2000)

[5] M. S. Knebelman, Conformal geometry of generalized metric spaces, Proc.nat.Acad. Sci.USA..15(1929) pp. 33-41 and 376-379.

[6] M. Matsumoto, On C - reducible Finsler spaces, Tensor, N.S.. 24(1972) pp. 29-37.

[7] M. Matsumoto, On Finsler spaces with Randers metric and special forms of important tensors, J. Math. Kyoto Univ.. 14(1974) pp. 477-498.

[8] R. Miron, General Randers space, Lagrange and Finsler geometry, Ed. by P.L. Antonelli and R.Miron..76(1996) pp.123-140.

[9] G. Randers, On the asymmetrical metric in the four- space of general relativity, Phys. Rev..2(1941)59 pp. 195-199.

[10] C. Shibata, On invariant tensors of β - change of Finsler metrics, J. Math. Kyoto Univ.. 24(1984) pp. 163-188.

[11] C. Shibata and M. Asuma, C-conformal invariant and tensors of Finsler metrics, Tensor, N.S.. 52(1993) pp. 76-81.

[12] C. Shibata and H. Shimada and M. Azuma and H. Yasda, On Finsler spaces with Randers metric, Tensor, N.S..31(1977) pp. 219-226.

[13] A. A. Tamim and N. L. Youssef, On generalized Randers manifold, Algebras, Groups and geometrys..16(1999) pp.115-126.